# **Re-managing the Flow**

The major rivers and streams of the Sacramento Valley provide essential pathways for spawning salmon and steelhead. Flow agreements to benefit these fish are on every major watercourse in the Sacramento Valley.

**Trinity** and **Shasta Lakes** are important sources of cold water storage. Timing the release of this cold water into the rivers is vital if spawning fish are to thrive.

Nhiskeytowr

#### Clear Creek 🕞

In May and June, water is pulsed into Clear Creek to attract Spring-run salmon from the Sacramento River. From June through October, water released from Whiskeytown Reservoir keeps water temperatures cool.

#### Sacramento River below Keswick Dam

In 1960, flow objectives were established for the protection of fish and wildlife. In 1990 and 1991 this policy was modified requiring more cold water when warmer temperatures would be harmful to fish.

#### Sacramento River at Wilkins Slough O

The Rivers and Harbors Act of 1935 mandated a specific flow rate at Wilkins Slough be maintained. The primary goals at that time were navigation and flood control. In 1992, Congress made protection of fish and wildlife a secondary goal and this requirement was updated in 2009.



For more details visit www.norcalwater.org/ efficient-water-management/instream-flows/ Sutter Buttes

Shasta Lake



Various flow agreements benefit spring run salmon.

## Feather River

A water quality certification adopted in 2010 provides for specific flow and temperature requirements to accommodate spawning salmon and steelhead.

> lew Bullards Bal Reservior

### Yuba River 🖒

In 2008, the Yuba River Accord increased the streamflow requirements over previous levels, which benefits fish while insuring sufficient water supplies for irrigation and municipal uses.

Folsom Lake

#### American River below Nimbus Dam

In 2000, the Flow Management Standard was developed, which established minimum flow standards to improve the conditions for fall-run Chinook salmon and steelhead. Additionally, releases are adjusted to maintain sufficiently low water temperatures for steelhead rearing in summer and Chinook spawning in the fall.